

## Reminders

- Final Project Proposal Due tonight
  - 10% of your final project grade!
- HW 7 due Monday (skip Question 4 about the survey)
- Lab 9 this week
  - Last lab assignment of the semester!!
  - Thanksgiving next week, then final project consultations the week after
- I will not be having Wednesday office hours next week and the week after

## Last Topic: Prediction

- Today: Correlation and Linear Regression
- Monday, Nov 24: Least Squares and Residuals HW 7 due
- Wednesday, Nov 26: Holiday!
- Monday, Dec 1: Regression Inference 

  → HW 8 due

  Progress Report due
- Wednesday, Dec 3: Special Topics (Data Ethics)
  - Final Project Consultations during Lab
- Monday, Dec 8: Special Topics (Data Privacy) HW 9 due

# Final Project

- Report is a written report
  - Should be in paragraph form... Do not simply list bullet points!
- Progress report we are only checking the written report
  - You do not need to submit your notebooks

Update: Progress report is checking written report AND code

- Final Project Report you will submit the completed written report and your notebook
  - Due on the Friday Dec 12 at 11:59pm
  - No late days for the final report. Submit what you have by the deadline


# Correlation

## Prediction

- Guessing the future based on data
- To predict the value of a variable:
  - Identify (measurable) attributes that are associated with that variable
  - Describe the relation between the attributes and the variable you want to predict
  - Use the relation to predict the value of a variable

#### **Trend**

- Positive association
- Negative association
- Pattern




**Positive Linear Association** 


#### Any discernible "shape" in the scatter

- Linear
- Non-linear

#### **Trend**

- Positive association
- Negative association
- Pattern





#### Any discernible "shape" in the scatter


- Linear
- Non-linear


#### **Trend**


- Positive association
- Negative association
- Pattern



- Linear
- Non-linear









#### **Trend**


- Positive association
- Negative association
- Pattern




- Linear
- Non-linear









## Correlation Coefficient r

- A metric for the strength of the linear relationship between two variables
  - How clustered values are in a scatter plot around a straight line
- r is always between -1 and 1 ( $-1 \le r \le 1$ )
  - r = 1: perfect correlation (straight line) sloping upward
  - r = -1: perfect correlation (straight line) sloping down
- r = 0: no linear association (*uncorrelated*)

# Notebook demo: Correlation Plots

## Notes about using r


- r measures linear association
- Association does not imply causation
  - Two variables may be correlated, but one may not cause the other
  - Non-linearity and outliers can affect correlation
- Correlations based on aggregated data (ecological correlations) can be misleading
  - Correlations between individuals may be lower than correlation between averages of groups

## Questions

True or false?

1. If the correlation coefficient of x and y is 0, then knowing one cannot help us predict the other

# Non-linear correlations may have $r \approx 0$



x and y are highly correlated, but not linearly


$$r \approx 0$$

## Questions

#### True or false?

- 1. If the correlation coefficient of x and y is 0, then knowing one cannot help us predict the other
- 2. If x and y have a correlation coefficient of 1, then one must cause the other

# Association $\neq$ causation



## Questions

#### True or false?

- 1. If the correlation coefficient of x and y is 0, then knowing one cannot help us predict the other
- 2. If x and y have a correlation coefficient of 1, then one must cause the other
- 3. If x and y have a correlation coefficient of -0.8, they have a negative association


# Prediction

## Guess the future

- Based on incomplete information
- One way of making predictions:
  - To predict the outcome for an individual, find others who are like that individual and whose outcomes you know. Use those outcomes as the basis of your prediction

# Example: Galton's Heights


Goal: Predict the height of a new child based on that child's midparent height (average parent height)



## Example: Galton's Heights

Goal: Predict the height of a new child based on that child's midparent height (average parent height)


- How could we predict the child's height given a mid parent height of 68 inches?



# Example: Galton's Heights

Goal: Predict the height of a new child based on that child's midparent height (average parent height)

- How could we predict the child's height given a mid parent height of 68 inches?
- Idea: Use the average height of the children of families whose midparent height is close to 68 inches




# Notebook Demo: Child's Height


## Graph of Average

- For each x value, the prediction is the average of the y values in its nearby group
  - The graph of these predictions is the graph of averages
- If the association between x and y is linear, then points in the graph of averages tend to fall on a line
  - This line is called the regression line

# Graph of Average




# Graph of Average



# Nearest Neighbor Regression

- A method for predicting a numerical y given a value of x:
  - Identify the group of points
     where the values of x are close
     to the given value
  - The prediction is the average of the y values for the group



# Calculating r

## Formalula for r

- The correlation coefficient (r) is the average product of  $\boldsymbol{x}$  in standard units and  $\boldsymbol{y}$  in standard units
  - To determine r, we first convert our values in x & y to standard units
  - We'll denote  $x_{su}$  and  $y_{su}$ , respectively

$$y_{su} = r \times x_{su}$$


This is known as the linear regression line

## Standard Units

- The quantity z (from "average  $\pm z$  SDs" in Chebychev's inequality) measures **standard units** 
  - Standard units is the number of standard deviations away from the average
- To convert a value (v) to standard units, compare the deviation from the average ( $\mu$ ) with the standard deviation (SD):

$$z = \frac{v - \mu}{SD}$$

## Converting to Standard Units



## Correlation Coefficient r

- r is the average of the product of two variables, when both variables are measured in standard units
- What this means for us:
  - r is not affected by changing the units of the measurement of the data
  - r will be the same regardless of which variable is plotted on the x- and y- axes

# Regression Line: Slope & Intercept


## Regression Line

Lines generally can be expressed by

$$y = slope \times x + intercept$$

When in standard units, this is equivalent to

$$y_{\rm su} = r \times x_{\rm su}$$





# Regression Line: Original Units

$$y_{\rm su} = r \times x_{\rm su}$$

$$\frac{\text{estimate of } y - \text{avg}(y)}{\text{SD of } y} = r \times \frac{x - \text{avg}(x)}{\text{SD of } x}$$


# Regression Line: Original Units





# Regression Line: Original Units

#### **Original Units**



estimate of  $y = \text{slope} \times x + \text{intercept}$ 

slope = 
$$r \times \frac{\text{SD of } y}{\text{SD of } x}$$

intercept = avg(y) - slope  $\times avg(x)$ 

# Notebook Demo: Regression Line for Height

### Next time

- Today: Correlation and Linear Regression

- Monday, Nov 24: Least Squares and Residuals - HW 7 due

- Wednesday, Nov 26: Holiday!
- Monday, Dec 1: Regression Inference 

  → HW 8 due

  Progress Report due
- Wednesday, Dec 3: Special Topics (Data Ethics)
  - Final Project Consultations during Lab
- Monday, Dec 8: Special Topics (Data Privacy) HW 9 due